



## Algebra 2/Trig. Cheat Sheet


### Unit Circle:



### Special Triangles:



### Trigonometric Functions: (SOH CAH TOA) (Only work on right triangles.)



$$\sin(\theta) = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\cos(\theta) = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\tan(\theta) = \frac{\text{opposite}}{\text{adjacent}}$$

### Trigonometric Inverses/Identities:

$$\csc(\theta) = \frac{1}{\sin(\theta)} \quad \sec(\theta) = \frac{1}{\cos(\theta)}$$

$$\cot(\theta) = \frac{\cos(\theta)}{\sin(\theta)} \quad \cot(\theta) = \frac{1}{\tan(\theta)}$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$$

### Co-function Identities:

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos(\theta)$$

$$\csc\left(\frac{\pi}{2} - \theta\right) = \sec(\theta)$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \cot(\theta)$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin\theta$$

$$\sec\left(\frac{\pi}{2} - \theta\right) = \csc(\theta)$$

$$\cot\left(\frac{\pi}{2} - \theta\right) = \tan(\theta)$$

### Pythagorean Identities:

$$\sin^2\theta + \cos^2\theta = 1$$

$$1 + \tan^2\theta = \sec^2\theta$$

$$1 + \cot^2\theta = \csc^2\theta$$

### Trig. Proofs:

$$\tan\theta = \sin\theta \sec\theta$$

$$\tan\theta = \sin\theta \times \frac{1}{\cos\theta}$$

$$\tan\theta = \frac{\sin\theta}{\cos\theta}$$

$$\tan\theta = \tan\theta$$

### Converting Degrees to Radians:

$$60^\circ \times \frac{\pi}{180} = \frac{\pi}{3}$$

### Converting Radians to Degrees:

$$\frac{\pi}{3} \times \frac{180}{\pi} = 60^\circ$$

**Distance Formula:**

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

**Law of Cosines:**  $c^2 = a^2 + b^2 - 2ab\cos(c)$ **Product of Roots:**  $\frac{c}{a}$ **Sum of Roots:**  $-\frac{b}{a}$ **Law of Sines:**

(This will be included on reference table)

$$\frac{\sin(A)}{a} = \frac{\sin(B)}{b} = \frac{\sin(C)}{c}$$

OR

$$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$$

**Solving Trig. Functions Algebraically:**Solve for  $\theta$ , in the interval  $-2\pi < \theta < 2\pi$ 

$$\sin^2\theta - 1 = 0$$

$$(\sin\theta + 1)(\sin\theta - 1) = 0$$

$$\sin\theta + 1 = 0$$

$$\sin\theta = -1$$

$$\sin^{-1}(-1) =$$

$$\theta = -90^\circ, 270^\circ$$

$$\sin\theta - 1 = 0$$

$$\sin\theta = 1$$

$$\sin^{-1}(1) =$$



$$\theta = -270^\circ, 90^\circ$$

$$\theta = -270^\circ, -90^\circ, 90^\circ, 270^\circ$$

**Logarithms are Inverses of Exponential Equations:**

$$y = \log_{10}(x)$$

$$y = 10^x$$

**Logarithm Rules:**

$$\log_{10}(100) = 2 \longrightarrow 10^2 = 100$$

$$\log(ab) = \log(a) + \log(b)$$

$$\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$$

$$\log a^b = b \cdot \log(a)$$

$$\ln_e(x) = e^x$$

**To Change Log Base:**

$$\log_a(x) = \frac{\log_b(x)}{\log_b(a)}$$

**Logarithmic Equations:**

$$\log_4(x) + \log_4(x + 6) = 2$$

$$\log_4(x(x + 6)) = 2$$

$$\log_4(x^2 + 6x) = 2$$

$$4^2 = x^2 + 6x$$

$$x^2 + 6x - 16 = 0$$

$$(x + 8)(x - 2) = 0$$

$$x + 8 = 0 \quad | \quad x - 2 = 0$$

$$x = -8 \quad | \quad x = 2$$

$$x = -8, 2$$

**Radical Equations:**

$$10\sqrt{x+4} = 100$$

$$\frac{10}{10} \sqrt{x+4} = \frac{100}{10}$$

$$\sqrt{x+4} = 10$$

$$(\sqrt{x+4})^2 = 10^2$$

$$x + 4 = 100$$

$$x = 96$$

**Simplifying Radical Expressions:**

$$\sqrt{40} = \sqrt{4 \cdot 10} = \sqrt{4}\sqrt{10} = 2\sqrt{10}$$

$$\sqrt[4]{256m^4n^8} = 4mn^2$$

$$25^{-\frac{1}{2}} = \frac{1}{\sqrt{25}} = \frac{1}{5}$$

**Rationalizing Radical in denominator:**

$$\frac{1}{\sqrt{x}} \times \frac{\sqrt{x}}{\sqrt{x}} = \frac{\sqrt{x}}{x}$$

**Radical Rules:**

$$\sqrt[n]{x} = x^{\frac{1}{n}}$$

$$x^{\frac{m}{n}} = (\sqrt[n]{x})^m$$

---

**Imaginary Numbers:**

$i^0 = 1$   
 $i^1 = i$   
 $i^2 = -1$   
 $i^3 = -i$   
 $a + bi$   
 Given  $i^{64}$   
 divide by 4 using  
 long division, get  
 remainder 0, then  
 use as exponent.  
 $i^{64} \rightarrow i^0 = 1$

**Sequences:**

Arithmetic Sequence: (add or subtract)

$$a_n = a_1 + (n - 1)d$$

Geometric Sequence: (multiply or divide)

$$a_n = a_1 \cdot r^{n-1}$$


---

**Binomial Theorem:**

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

**Equation of a Circle:**

$$(x - 4)^2 + (y + 2)^2 = 16$$

Center= (4, -2)

Radius= $\sqrt{16} = 4$ 

---

**Probability:**Binomial Probability=  ${}_n C_r \cdot p^r \cdot q^{n-r}$ Dependent Events=  $P(A) \cdot P\left(\frac{B}{A}\right)$ 

P (A and B)

Permutation=  ${}_n P_r = \frac{n!}{(n-r)!}$   
(Order matters)Independent Events= $P(A) \cdot P(B)$ 

P (A and B)

Combinations=  ${}_n C_r = \frac{n!}{r!(n-r)!}$   
(Order doesn't matter)Not mutually Exclusive=  $P(A) + P(B) - P(A \text{ or } B)$ 

P (A or B)

Conditional=  $\frac{P(A \cap B)}{P(A)} = P(B|A)$ Mutually Exclusive= $P(A) + P(B)$ P (A or B)

---

**Population Mean ( $\mu$ ):**

$$\mu = \frac{1}{n} \sum_{i=1}^n x_i$$

**Population Standard Deviation ( $\sigma$ ):**

$$\sigma = \sqrt{\frac{\sum_{i=1}^n (x - \mu)^2}{N}}$$

$$z = \frac{x - \mu}{\sigma}$$

**Sample Mean ( $\bar{x}$ ):**

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$



**Sample Standard Deviation (S):**

$$S = \sqrt{\frac{\sum_{i=1}^n (x - \bar{x})^2}{n - 1}}$$

**Margin of Error:**

$$MOE = z \times \frac{S}{\sqrt{n}}$$


---

**Venn Diagram:** $A \cup B$  $A \cap B$ 

**Absolute Value:** When solving for x create two equations. One with a positive answer, one with a negative. Then solve as normal. Ex:

$$|-125 + 2x| = 45$$

$$\begin{array}{l} -125 + 2x = 45 \\ x = 85 \end{array} \qquad \begin{array}{l} -125 + 2x = -45 \\ x = 40 \end{array}$$

**Factoring Methods to Know:**

- 1) Greatest Common Factor (GCF)
- 2) Quadratic Formula:  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
- 2) Completing the Square
- 3) Difference of Two Squares (DOTS)
- 4) Factor by Grouping

**Binomial Expansion:**

$$a^2 - b^2 = (a - b)(a + b)$$

$$a^3 + b^3 = (a + b)(a^2 + 2ab + b^2)$$

$$a^3 - b^3 = (a - b)(a^2 - 2ab + b^2)$$

**Discriminant Rules:**

$$b^2 - 4ac$$

$b^2 - 4ac > 0$	2 real roots
$b^2 - 4ac = 0$	1 real root
$b^2 - 4ac < 0$	2 imaginary roots

**Dividing Polynomials:**

$$\begin{array}{r}
 2x + 1 \overline{) 6x^3 + 5x^2 + 3x + 4} \\
 \underline{6x^3 + 3x^2} \phantom{+ 4} \\
 2x^2 + 3x + 4 \\
 \underline{2x^2 + x} \phantom{+ 4} \\
 \phantom{2x^2} 2x + 4 \\
 \underline{\phantom{2x^2} 2x + 1} \\
 \phantom{2x^2} \phantom{2x} 3
 \end{array}$$

R: 3

Answer:  $3x^2 + x + 1 + \frac{3}{2x + 1}$

**Standard Form of a Parabola:**

$$y = ax^2 + bx + c$$

Axis of Symmetry:  $-\frac{b}{2a}$

**Vertex Form of a Parabola:**

$$y = a(x - h)^2 + k$$

Vertex:  $(h, k)$

- $0 < a < 1$ , then parabola gets bigger
- $a > 0$ , then parabola gets narrow

**Remainder Theorem:** If polynomial  $f(x)$  is divided by  $(x - a)$ , then remainder is equal to  $f(a)$ . Ex:

If  $f(x) = 4x^2 + 6x + 15$  is divided by  $(x - 1)$

The remainder will be...

$$f(1) = 4(1)^2 + 6(1) + 15 = 25$$

**Simplifying Rational Expressions:**

$$\frac{9y^3 - y}{3y - 1} = \frac{y(9y^2 - 1)}{3y - 1} = \frac{y(3y - 1)(3y + 1)}{3y - 1} = y(3y + 1)$$

**Compound Interest Formula:**

$$A = P\left(1 + \frac{r}{n}\right)^{nt}$$

- $P$  = Principle
- $r$  = Interest rate
- $n$  = number of compoundings per year
- $t$  = Total number of years

**Exponential Equation:**

$$y = ab^x$$

**Exponential Growth:**

$$b > 1$$

**Exponential Decay:**

$$0 < b < 1$$

---

**Power Functions:**

$$y = kx^n$$

where,  $k$  and  $n$  are known number constants.


**Odd Power Functions:**  $y = kx^n$ 

where  $n$  is equal to any odd number

{1,3,5,7, ... ..}

Ex:

$$y = x^3$$


**Even Power Functions:**  $y = kx^n$ 

where  $n$  is equal to any even number

{2, 4, 6, 8, ... ..}

Ex:


$$y = x^2$$



---

**Focus and Directrix:**

A parabola is equi-distant between both focus and directrix.



To find an equation of parabola, given focus (2,1) and directrix  $y = -1$ :


$$\sqrt{(y - \text{directrix})^2} = \sqrt{(x - a)^2 + (y - b)^2}$$

(where  $a$  and  $b$  are from focus)

$$\sqrt{(y + 1)^2} = \sqrt{(x - 2)^2 + (y - 1)^2}$$

$$y = \frac{(x - 2)^2}{4}$$


**Transformations of Function  $f(x) = x^2$ , where  $C = 2$ :**



Function Transformation	What does it do to the graph?	Graph
$y = f(x) + C$	$C > 0$ moves up $C < 0$ moves down	<p><math>f(x) = x^2 + 2</math></p>
$y = f(x + C)$	$C > 0$ moves left $C < 0$ moves right	<p><math>f(x) = (x^2 + 2)</math></p>
$y = Cf(x)$	$C > 1$ moves closer to y-axis $0 < C < 1$ moves further from y-axis	<p><math>f(x) = 2x^2</math></p>
$y = -f(x)$	Reflection in the x-axis	<p><math>f(x) = -x^2</math></p>

$$y = f(-x)$$

*Reflection in the y - axis*



$$f(x) = -x^2$$